

Catch-up Immunization Schedule for Persons Aged 4 Months Through 18 Years Who Start Late or Who Are More Than 1 Month Behind—United States • 2010

The table below provides catch-up schedules and minimum intervals between doses for children whose vaccinations have been delayed. A vaccine series does not need to be restarted, regardless of the time that has elapsed between doses. Use the section appropriate for the child's age.

PERSONS AGED 4 MONTHS THROUGH 6 YEARS					
Vaccine	Minimum Age for Dose 1	Minimum Interval Between Doses			
		Dose 1 to Dose 2	Dose 2 to Dose 3	Dose 3 to Dose 4	Dose 4 to Dose 5
Hepatitis B ¹	Birth	4 weeks	8 weeks (and at least 16 weeks after first dose)		
Rotavirus ²	6 wks	4 weeks	4 weeks ²		
Diphtheria, Tetanus, Pertussis ³	6 wks	4 weeks	4 weeks	6 months	6 months ³
<i>Haemophilus influenzae</i> type b ⁴	6 wks	4 weeks if first dose administered at younger than age 12 months 8 weeks (as final dose) if first dose administered at age 12–14 months No further doses needed if first dose administered at age 15 months or older	4 weeks ⁴ if current age is younger than 12 months 8 weeks (as final dose)⁴ if current age is 12 months or older and first dose administered at younger than age 12 months and second dose administered at younger than 15 months No further doses needed if previous dose administered at age 15 months or older	8 weeks (as final dose) This dose only necessary for children aged 12 months through 59 months who received 3 doses before age 12 months	
Pneumococcal ⁵	6 wks	4 weeks if first dose administered at younger than age 12 months 8 weeks (as final dose for healthy children) if first dose administered at age 12 months or older or current age 24 through 59 months No further doses needed for healthy children if first dose administered at age 24 months or older	4 weeks if current age is younger than 12 months 8 weeks (as final dose for healthy children) if current age is 12 months or older No further doses needed for healthy children if previous dose administered at age 24 months or older	8 weeks (as final dose) This dose only necessary for children aged 12 months through 59 months who received 3 doses before age 12 months or for high-risk children who received 3 doses at any age	
Inactivated Poliovirus ⁶	6 wks	4 weeks	4 weeks	6 months	
Measles, Mumps, Rubella ⁷	12 mos	4 weeks			
Varicella ⁸	12 mos	3 months			
Hepatitis A ⁹	12 mos	6 months			
PERSONS AGED 7 THROUGH 18 YEARS					
Tetanus, Diphtheria/ Tetanus, Diphtheria, Pertussis ¹⁰	7 yrs ¹⁰	4 weeks	4 weeks if first dose administered at younger than age 12 months 6 months if first dose administered at 12 months or older	6 months if first dose administered at younger than age 12 months	
Human Papillomavirus ¹¹	9 yrs		Routine dosing intervals are recommended¹¹		
Hepatitis A ⁹	12 mos	6 months			
Hepatitis B ¹	Birth	4 weeks	8 weeks (and at least 16 weeks after first dose)	6 months	
Inactivated Poliovirus ⁶	6 wks	4 weeks	4 weeks	6 months	
Measles, Mumps, Rubella ⁷	12 mos	4 weeks			
Varicella ⁸	12 mos	3 months if person is younger than age 13 years 4 weeks if person is aged 13 years or older			

1. Hepatitis B vaccine (HepB).

- Administer the 3-dose series to those not previously vaccinated.
- A 2-dose series (separated by at least 4 months) of adult formulation Recombivax HB is licensed for children aged 11 through 15 years.

2. Rotavirus vaccine (RV).

- The maximum age for the first dose is 14 weeks, 6 days. Vaccination should not be initiated for infants aged 15 weeks, 0 days or older.
- The maximum age for the final dose in the series is 8 months, 0 days.
- If Rotarix was administered for the first and second doses, a third dose is not indicated.

3. Diphtheria and tetanus toxoids and acellular pertussis vaccine (DTaP).

- The fifth dose is not necessary if the fourth dose was administered at age 4 years or older.

4. *Haemophilus influenzae* type b conjugate vaccine (Hib).

- Hib vaccine is not generally recommended for persons aged 5 years or older. No efficacy data are available on which to base a recommendation concerning use of Hib vaccine for older children and adults. However, studies suggest good immunogenicity in persons who have sickle cell disease, leukemia or HIV infection, or who have had a splenectomy; administering 1 dose of Hib vaccine to these persons who have not previously received Hib vaccine is not contraindicated.
- If the first 2 doses were PRP-OMP (PedvaxHIB or Comvax) and administered at age 11 months or younger, the third (and final) dose should be administered at age 12 through 15 months and at least 8 weeks after the second dose.
- If the first dose was administered at age 7 through 11 months, administer the second dose at least 4 weeks later and a final dose at age 12 through 15 months.

5. Pneumococcal vaccine.

- Administer 1 dose of pneumococcal conjugate vaccine (PCV) to all healthy children aged 24 through 59 months who have not received at least 1 dose of PCV on or after age 12 months.
- For children aged 24 through 59 months with underlying medical conditions, administer 1 dose of PCV if 3 doses were received previously or administer 2 doses of PCV at least 8 weeks apart if fewer than 3 doses were received previously.
- Administer pneumococcal polysaccharide vaccine (PPSV) to children aged 2 years or older with certain underlying medical conditions, including a cochlear implant, at least 8 weeks after the last dose of PCV.

6. Inactivated poliovirus vaccine (IPV).

- The final dose in the series should be administered on or after the fourth birthday and at least 6 months following the previous dose.

- A fourth dose is not necessary if the third dose was administered at age 4 years or older and at least 6 months following the previous dose.
- In the first 6 months of life, minimum age and minimum intervals are only recommended if the person is at risk for imminent exposure to circulating poliovirus (i.e., travel to a polio-endemic region or during an outbreak).

7. Measles, mumps and rubella vaccine (MMR).

- Administer the second dose routinely at age 4 through 6 years. However, the second dose may be administered before age 4, provided at least 28 days have elapsed since the first dose.
- If not previously vaccinated, administer 2 doses with at least 28 days between doses.

8. Varicella vaccine.

- Administer the second dose routinely at age 4 through 6 years. However, the second dose may be administered before age 4, provided at least 3 months have elapsed since the first dose.
- For persons aged 12 months through 12 years, the minimum interval between doses is 3 months. However, if the second dose was administered at least 28 days after the first dose, it can be accepted as valid.
- For persons aged 13 years and older, the minimum interval between doses is 28 days.

9. Hepatitis A vaccine (HepA).

- HepA is recommended for children older than 23 months who live in areas where vaccination programs target older children who are at increased risk for infection, or for whom immunity against hepatitis A is desired.

10. Tetanus and diphtheria toxoids vaccine (Td) and tetanus and diphtheria toxoids and acellular pertussis vaccine (Tdap).

- Doses of DTaP are counted as part of the Td/Tdap series
- Tdap should be substituted for a single dose of Td in the catch-up series or as a booster for children aged 10 through 18 years; use Td for other doses.

11. Human papillomavirus vaccine (HPV).

- Administer the series to females at age 13 through 18 years if not previously vaccinated.
- Use recommended routine dosing intervals for series catch-up (i.e., the second and third doses should be administered at 1 to 2 and 6 months after the first dose). The minimum interval between the first and second doses is 4 weeks. The minimum interval between the second and third doses is 12 weeks, and the third dose should be administered at least 24 weeks after the first dose.

Information about reporting reactions after immunization is available at www.vaers.hhs.gov or by calling 800-822-7967. Suspected cases of vaccine-preventable diseases should be reported to the state or local health department. Additional information, including precautions and contraindications for immunization, is available from the National Center for Immunization and Respiratory Diseases at www.cdc.gov/vaccines or 800-CDC-INFO (800-232-4636)